



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Using this fact, a little more reduction leads to the result

$$R = \frac{(a+e-f)}{24V} \sqrt{[(e-c)^2-d^2][(c-a)^2-b^2][f^2-(a-e)^2]}, \quad (3)$$

where

$$144V^2 = a^2d^2(e^2+f^2-a^2) + b^2e^2(f^2+a^2-e^2) + c^2f^2(a^2+e^2-f^2) \\ - a^2(d^2-b^2)(d^2-c^2) - e^2(b^2-c^2)(b^2-d^2) - f^2(c^2-d^2)(c^2-b^2) - a^2e^2f^2. \quad (4)$$

Seven similar expressions can be found for the radii of spheres tangent to the four edges a, c, e, f , touching one or more of the edges produced beyond the tetrahedron.

Thirdly, applying the conditions (1) to (3), we can obtain the required answer in a symmetrical form. Now (3) is the radius of a sphere touching the four edges a, c, e, f . If this sphere touches also the edges b, d , that is, if $a+d = b+e = c+f$, then

$$R = \frac{1}{24V} \sqrt[3]{\frac{(a+b-c)(a-b+c)(-a+b+c)(b+d-f)}{(b-d+f)(-b+d+f)(c+d-e)(c-d+e)} \\ (-c+d+e)(a+e-f)(a-e+f)(-a+e+f)}, \quad (5)$$

where V has the value given in (4), which is already expressed in a symmetrical form. This is the required solution.

It may be added that, if A_1, A_2, A_3, A_4 are the areas of the four face-triangles, and r_1, r_2, r_3, r_4 the radii of the circles inscribed to these triangles, the above expression takes the simple form

$$R = \frac{2}{3V} \sqrt[3]{A_1A_2A_3A_4r_1r_2r_3r_4}. \quad (6)$$

If the tetrahedron is regular the expression reduces to

$$R = \frac{a}{2\sqrt{2}}.$$

Similarly the radii of the four spheres R_1, R_2, R_3, R_4 which touch three edges and the other three edges produced, are

$$R_1 = \frac{2}{3V} \sqrt[3]{A_1A_2A_3A_4r_{121}r_{31}r_{41}}; \quad R_2 = \frac{2}{3V} \sqrt[3]{A_1A_2A_3A_4r_{212}r_{32}r_{42}}; \\ R_3 = \frac{2}{3V} \sqrt[3]{A_1A_2A_3A_4r_{313}r_{23}r_{43}}; \quad R_4 = \frac{2}{3V} \sqrt[3]{A_1A_2A_3A_4r_{414}r_{34}r_{24}},$$

where r_{pq} is the radius of the circle escribed to the triangle A_p which touches the edge common to A_p and A_q and the other two edges of A_p produced.

If the tetrahedron is regular, $R_1 = R_2 = R_3 = R_4 = \frac{3a}{2\sqrt{2}}.$

CALCULUS.

358. Proposed by C. N. SCHMALL, New York City.

About a given circle circumscribe the smallest parabola.


SOLUTION BY HORACE OLSON, Chicago, Illinois.

We assume that by the smallest parabola is meant the smallest segment of a parabola having its bounding ordinate tangent to the given circle.

Let $PFRD$ be the given circle whose center C is the origin of rectangular coördinates. Then, without loss of generality, the parabola may be assumed to have its axis on the axis of abscissas AD .

Let EAG be the required parabola. Then, if r is the radius of the circle, we have $x^2 + y^2 = r^2$ for the circle, and $y^2 = 2p(x + k)$ for the parabola, where p is the distance from the focus to the directrix and k is the distance AC .

Since P is the point of tangency, solving these two simultaneous equations subject to the condition of tangency, we have $r^2 - 2kp + p^2 = 0$.



Hence, $k = p/2 + r^2/2p$.

Then

$$AD = AC + CD = \frac{p}{2} + \frac{r^2}{2p} + r = \frac{(r + p)^2}{2p},$$

and

$$ED = \sqrt{2p(AD)} = (r + p).$$

Area of $EAG = a = \frac{2}{3}AD \cdot EG = \frac{4}{3}(r + p)^2/2p \cdot (r + p) = \frac{2}{3}(r + p)^3/p$.

Equating to zero the derivative of a with respect p , we have

$$\frac{da}{dp} = \frac{2}{3} \cdot \frac{3p(r + p)^2 - (r + p)^3}{p^2} = 0;$$

whence

$$(p + r)^2 = 0 \quad \text{and} \quad 2p - r = 0.$$

Hence, $p = -r$ or $p = \frac{1}{2}r$. The value $p = -r$ gives neither a maximum nor a minimum. The value $p = \frac{1}{2}r$ gives a minimum, and the equation of the corresponding parabola is $y^2 = r(x + 5r)/4$.

A similar solution was received from the PROPOSER.

359. Proposed by W. D. CAIRNS, Oberlin College.

Examine for maxima and minima

$$f(x) = e^{-cx}(1 + \cos x) \quad (c > 0)$$